viſion ceaſes, and the ſecond leaf terminates in diſtinct, ſharply ſerrated lobes.

The ſeed-veſſels are placed in round dots, on the under ſide of the lobes; two or three on the ſmaller, five or ſix on the larger; they are of a brown colour, and diſtinct. The whole plant is larger, firmer, more robuſt, and of a ſtronger green than the Polypodium fragile.

The Polypodium fragile in ſome of its ſtates, ſee part 1, page 51, ſo nearly reſembles the Polypodium rhœticum, that others, as well as myſelf, have been deceived by it. Hallar, Wais, &c. ſeem to doubt its exiſtence; and it appears from their deſcriptions, that they had not examined the true Polypodium rhœticum; no more had I, otherwiſe ſo obvious a character as the inequality of ſize in the third leaves, on the oppoſite ſides of the rib of the ſecond leaf, could not have been diſregarded.

In my plate of varieties, *Part* 1, *Tab.* 2, thoſe of the Polypodium fragile not being ſufficiently expreſſed, at the inſtance of ſeveral friends, I have added *TAB. XLVI*; in which I have carefully copied theſe varieties. For on the two ſtems given in the plate, by varying the ſecond leaves, as I found them in ſeparate ſpecimens, I have repreſented all thoſe varieties which have been ſuppoſed moſt to reſemble the Polypodium rhœticum.

Specimens of Polypodium rhœticum and Acroſtichum thelypteris were ſent me by my friend Dickſon, author of the *Plantarum Cryptogamicarum Britanniæ*; the firſt he gathered in *Scotland*, the laſt, he informs me, grows in plenty about *Norwich*.

ERRATA IN THE FIRST PART.

INTRODUCTION, p. 6, l. 17, *for* ſupine *r.* prone.—P. 16, l. 3. *r.* Anglica.——*In the* BOOK, p. 5, l. 3. *r.* and when the ſeeds are diſcharged.—P. 40, l. 2, 3, *eraſe* Ger. Em. & Park.—P. 41, l. 3, tab. 22.—P. 41, l. 3, *eraſe* from this, *and the reſt of the ſentence.*—P. 51, l. 15, *eraſe* this is, *and the reſt of the ſentence*;—*ſame line, after* delicate, *add* tab. 46.—*Throughout the Book, for* Polipodium *r.* Polypodium.